

Synthesis, Crystal Structures of S- and N-Benzyl Derivatives Of 5-phenyl-1,3,4-oxadiazol-2-thione

A.A. Ziyaev¹, U.S. Makhmudov^{1*}, M.A. Ziyaeva², B. Tashkhodjaev¹

¹S.Yunusov Institute of the Chemistry of Plant Substances,
Academy of Sciences of Republic of Uzbekistan, Uzbekistan

²Tashkent technical university, Tashkent, Uzbekistan

Abstract: Benzylation reaction is carried out 5-phenyl-1,3,4-oxadiazol-2-thione with benzyl chloride in acetone and HMPA. In the acetone was obtained exclusively S-benzyl derivative and in the case of HMPA - isomeric mixture of S- and N-benzyl derivatives were obtained, individually isolated, characterized, and their structure was proved by spectral methods. By slow evaporation of solvents at room temperature were obtained two polymorphic crystals of 5-phenyl-1,3,4-oxadiazol-2-thione from acetone **1** and ethanol **2**. Crystals of 2-benzylthio-5-phenyl-1,3,4-oxadiazole **3** and 3-benzyl-5-phenyl-1,3,4-oxadiazol-2-thione **4** obtained in similar conditions have structures without the inclusion of solvent molecules.

Keywords: Five-membered heterocycles; 5-phenyl-1,3,4-oxadiazol-2-thione; 2-benzylthio-5-phenyl-1,3,4-oxadiazole; 3-benzyl-5-phenyl-1,3,4-oxadiazol-2-thione; X-ray analysis; Crystal structure

Received: January 29, 2017; **Accepted:** February 18, 2017; **Published:** March 15, 2017

Competing Interests: The authors have declared that no competing interests exist.

Copyright: 2016 U.S. Makhmudov, *et al.* This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

***Correspondence to:** U.S. Makhmudov, S.Yunusov Institute of the Chemistry of Plant Substances, Academy of Sciences of Republic of Uzbekistan, Uzbekistan

Email: Earl-Sharp@rambler.ru

Introduction

5-Substituted-1,3,4-oxadiazol-2-thiones interesting representatives of one five-membered heterocycles. Numerous compounds of this class show practical interest for researchers having a broad spectrum of biological activity, such as pesticide, antibiotic, anticancer, antihypertensive, antidepressant, analgesic, anti-inflammatory and others [1-4]. The presence in its molecule ambifunctional thioamide NH-C=S and other functional groups gives oxadiazolthiones to exhibit multiple reactivity whereby depending on the reaction conditions and the nature of the reagents can be prepared isomeric (S-, N-) products.

In this paper the task was to continue previously initiated a study on the alkylation reactions oxadiazol-2-thiones [5, 6], namely, to synthesize isomeric S- and N-benzyl derivatives of N-5-phenyl-1,3,4-oxadiazol-2-thione, prove their structure and study crystal structures (Fig. 1).

Materials and Methods

UV spectra were recorded on a spectrometer EPS-3T Hitachi in ethanol. ^1H NMR spectra were recorded in CDCl_3 on Varian 400-MR spectrometer operating accordingly at 400 MHz. Hexamethyldisiloxane (HMDSO) was used as internal standard, chemical shift of ^1H was recorded in ppm. Melting point were measured on a Boetius and MEL-TEMP apparatus manufactured by Barnstead International (USA) and were uncorrected.

The reactionary process was monitored by TLC on ALUGRAM Xtra (Germany) precoated aluminum plates using $\text{CHCl}_3/\text{EtOH}$ (24:1) solvent system and developed plates were visualized under UV lamp and/or iodine tank where necessary.

The unit cell parameters of the crystals are determined and refined with CCD Xcalibur Ruby diffractometer (Oxford Diffraction) using $\text{CuK}\alpha$ -radiation, graphite monochromator ($T=293\text{ K}$). The three-dimensional set of reflections received at the appropriate diffractometer.

Results

Synthesis

Reaction 5-phenyl-1,3,4-oxadiazol-2-thione with benzyl chloride in acetone. The reaction in equimolar ratios of thione **1**, benzyl chloride and K_2CO_3 was performed by refluxing in dry acetone for 5 hours. Acetone was then removed from the mixture, residue was washed with water, then with alkali solution to remove residues thione and again with water until neutral pH. The resulting product **3** was dried and recrystallized from ethanol.

2-benzylthio-5-phenyl-1,3,4-oxadiazol 3: 95% yield, m.p.97-980°C, $R_f=0,71$ (chloroform-ethanol, 24: 1), UV spectrum, λ_{max} , 275nm; ^1H NMR spectrum (CDCl_3), δ , ppm, 4.50 (2H, s, S-CH₂), 7.25-8.20 (10H, m, Ar-H).

Reacting 5-phenyl-1,3,4-oxadiazol-2-thione with benzyl chloride in hexamethylphosphoric triamide (HMPA).

The reaction was performed by stirring equimolar amounts of thione **1**, benzyl chloride and K_2CO_3 in HMPA at a temperature of water bath (70-80 °C) for 5 hours. The reaction mixture was poured

into ice water, the precipitate was filtered, washed with an alkali solution and several times with cold water. TLC analysis showed two spots, i.e. presence S- and N-benzyl derivatives, which overall yield was 94%. Isolation of individual isomers was carried out on a silica gel column Chemapol 250/400 mm, eluent hexane-chloroform in a ratio of 5:1, 3:1, 1:1.

3-benzyl-5-phenyl-1,3,4-oxadiazol-2-thione 4: yield 3%, m.p.136-137⁰C, $R_f=0,88$ (chloroform-ethanol, 24:1), UV spectrum, λ_{max} , 302nm; ¹H NMR spectrum (CDCl₃), δ , ppm, 5.25 (2H, s, N-CH₂), 7.24-8.15 (10H, m, Ar-H).

Crystal growth

Single crystals of 5-phenyl-1,3,4-oxadiazol-2-thione were grown from solutions of acetone **1** and ethanol **2** by slow evaporation at room temperature. Crystals of 2-benzylthio-5-phenyl-1,3,4-oxadiazole (S-isomer) **3** and 3-benzyl-5-phenyl-1,3,4-oxadiazol-2-thione (N-isomer) **4** were grown from ethanol and acetone by slow evaporation of the solvent at room temperature.




Fig. 1 Schematic formula and numbering of the atoms in the structures **1,2,3** and **4**.

X-ray Structure Determination

The unit cell parameters of the crystals were determined and refined with CCD Xcalibur Ruby diffractometer (Oxford Diffraction) using CuK α -radiation, graphite monochromator (T=293 K). The three-dimensional set of reflections received at the appropriate diffractometer. The amendment was introduced to the absorption by Multi-scan [7]. Table 1 shows the main parameters of the X-ray diffraction experiments and refinement calculations of the structures **1,2,3** and **4**.

The structures were solved by direct methods using SHELXS-2014 and refined using SHELXL-2014 programs [8]. All non-hydrogen atoms were refined by anisotropic full-matrix least-squares methods (over F^2). Positions of H atoms were found geometrically and refined with fixed isotropic thermal parameters $U_{iso} = nU_{eq}$, where $n = 1.5$ for methyl hydrogens and 1.2 for others and U_{eq} is the equivalent isotropic thermal parameter of the corresponding C atoms.

Materials X-ray diffraction as a CIF file deposited at the Cambridge center of crystal data (CCDC), from which can be obtained free on request at the following link: www.ccdc.cam.ac.uk/data_request/cif

Table 1 Basic crystallographic parameters and characteristics of the X-ray diffraction for structures **1,2,3** and **4**.


Compound	1	2	3	4
Molecular formula	C ₈ H ₆ N ₂ OS	C ₈ H ₆ N ₂ OS	C ₁₅ H ₁₂ N ₂ OS	C ₁₅ H ₁₂ N ₂ OS
Mr	178.21	178.21	268.33	268.33
Crystal symmetry	Orthorhombic	Monoclinic	Orthorhombic	Monoclinic
Space group	Pbca	P2 ₁ /c	Pbca	P2 ₁ /n
Z	8	4	8	4
a, Å	10.017(2)	5.5767(4)	12.6101(6)	11.4540(9)
b, Å	8.6679(17)	19.9882(11)	8.3582(4)	4.6674(3)
c, Å	17.634(4)	7.2282(6)	25.8399(14)	25.078(2)
α, °	90	90	90	90
β, °	90	100.463(7)	90	101.617(8)
γ, °	90	90	90	90
V, Å ³	1531.1(5)	792.32(10)	2723.5(2)	1313.21(17)
ρ, g/cm ³	1.546	1.494	1.309	1.357
Crystal dimension (mm)	0.35x0.48x0.55	0.40x0.45x0.55	0.35x0.45x0.50	0.40x0.45x0.50
Range scanned, 2θ °	4.432≤θ≤75.799°	6.607≤θ≤72.807°	5.498≤θ≤75.318°	3.596≤θ≤75.691°
μ _{exp} (cm ⁻¹)	3.314	3.202	2.049	2.125
No. reflection collected	1580	1583	2835	2698
No. reflection with I>2σ(I)	1296	920	1910	1909
R ₁ (I>2σ(I) and total)	0.0614 (0.0737)	0.0624 (0.1065)	0.0420 (0.0703)	0.0593 (0.0863)
wR ₂	0.1740 (0.1985)	0.1507 (0.1842)	0.1269 (0.1597)	0.1565 (0.1735)
GOOF	0.727	0.988	0.975	1.103
Largest diff. peak and hole (e Å ⁻³)	0.673 and -0.644	0.404 and -0.251	0.192 and -0.158	0.312 and -0.228
CCDC	1530177	1530178	1530179	1530180

Discussion

The reaction of equimolar ratios of thione, benzyl chloride and K₂CO₃ performed by refluxing in dry acetone. The result obtained and identified with high (95%) yield of the product solely S-2-benzylthio-5-phenyl-1,3,4-oxadiazol **3**. Detection of possible N-benzyl derivative was failed by thin layer chromatography (TLC) or spectroscopic (UV, ¹H-NMR) methods, that shows, in these conditions the reaction occurs with the formation of only one isomer.

A small amount (3%) of N-benzyl product was allocated during the reaction in a polar solvent - HMPA. As it is known, increasing polarity aprotic solvent promotes the alkylation of the site with the highest electron density. The reason for this is the ability of these solvents to solvate cations, in the result coordination with ambident anion is weakened. This facilitates the reaction by an atom having the highest electron density, which is the nitrogen atom of ambifunctional system of oxadiazol-2-thiones. Individual S-

and N-benzyl derivatives were isolated by column chromatography and they differ in the R_f values of TLC (0,71 and 0,88 for N- and S- isomers respectively). The structures were proven by UV and ^1H NMR spectroscopy. Thus in the UV spectrum of S-derivative **3** observes the absorption at 275 nm, in the case of N- benzyl derivative **4** at 302 nm. Also chemical shifts of isomers are clearly different and appear as a singlet for protons of the methylene group in S- CH_2 at 4.50 ppm, and for N- CH_2 at 5.25 ppm. The signals of the protons of aromatic rings in the form of a complex signal are in the range 7.24-8.20 ppm.



We have previously found that N-(3-methylthio and N-(3-ethylthio-5-atsetoamido-1,2,4-thiadiazole-2-thione) derivatives of the alkaloid cytosine exhibit clathratogen property including different solvent molecules in their crystal lattices [9-12]. Although the alkaloid cytosine [13] and other derivatives [14, 15] (for example, acyl or benzyl derivatives of cytosine [16, 17]) have not such properties. For this reason, was interesting study of the crystal structures of related oxadiazoles which grown from different solvents.

On this basis, it was of interest to study solvate formation and polymorphism of oxadiazol-2-thiones - related to 1,2,4-thiadiazole-2-thiones. For this purpose were grown visually distinguished two type crystals of 5-phenyl-1,3,4-oxadiazol-2-thione from solutions of acetone **1** and ethanol **2**. Further study of crystals by X-ray diffraction, it was found that they are two different polymorphic modifications of compounds differing packing factor and not containing a solvent molecule (Fig.2).

Further, the crystal structures of two isomeric S- and N-alkyl derivatives of 5-phenyl-1,3,4-oxadiazol-2-thione were studied. Crystals of 2-benzylthio-5-phenyl-1,3,4-oxadiazole (S-isomer) **3** and 3-benzyl-5-phenyl-1,3,4-oxadiazol-2-thione (N-isomer) **4** were grown from ethanol and acetone by slow evaporation of the solvent at room temperature. By determination of the melting point and the unit cell parameters was found that in these solvents grown crystals without the inclusion of solvent molecules (Fig.2).

Five membered heterocyclic ring in molecular structures **1**, **2**, **3** and **4** comprise a flat (within ± 0.0015 , ± 0.0073 , ± 0.0012 , and ± 0.0038 Å respectively). The location of phenyl ring relative to the plane of the oxadiazole nucleus charactering with value of torsion angle O1-C5-C6-C7 (12.47, -3.38, -5.76 and 7.01 ° for **1**, **2**, **3** and **4**, respectively). Namely the difference in the values of the torsion angle O1-C5-C6-C7 leads to the formation of different space groups and polymorphs in **1** and **2**.




Fig.2 Packing structures **1-4**.

In the structures **1,2,3** and **4** was observed weak $\pi \dots \pi$ interaction between the five and six membered aromatic rings (atoms O1, C2, N3, N4, C5 and C6, C7, C8, C9, C10, C11) of molecules related by symmetry $1/2-x, -1/2+y, z; x, 3/2-y, -1/2+z; 1/2-x, -1/2+y, z; x, -1+y, z$ and with centroid-centroid distance of 3.509 (2), 3.569(3), 3.7270(15), 3.628(2) Å respectively [18].

Thus, were synthesized and proven structures of S- and N-isomers of benzyl-5-phenyl-1,3,4-oxadiazol-2-thione. In the single crystals of isomers **3** and **4** grown from different solvents, unlike crystals of **1** and **2**, is not observed the formation of polymorphs.

In conclusion, it should be noted that the crystals grown from different solvents 5-phenyl-1,3,4-oxadiazole derivatives do not contain solvent molecules and they are not clathrogen.

References

1. Rakesh RS, Prabhakar Y. Shirodkar Oxadiazole: A biologically important heterocycle. *Der Pharma Chemica*. 2009, 1:130-140

2. Kharb R, Kaur R, Sharma AK. Vistas on antimicrobial potential of novel oxadiazole derivatives in modern medicinal chemistry. *European Journal of Biomedical and Pharmaceutical Sciences*. 2014,1:401-420
3. Kumar B, Kumar A, Beheraand AK, Raj V. Latest update on pharmacological activities of 1,3,4-oxadiazole derivatives. *J Cell Sci Ther*. 2016, 7:233-239
4. Jagadeesh PD, Shivarama HB, Nalilu SK, Laxmana K, Kumara Ch. Synthesis and antimicrobial evaluation of some new mannich bases bearing 1,3,4-oxadiazoline ring system. *International Journal of Advanced Research in Chemical Science*. 2015, 2:7-14
5. Ziyaev AA, Tozhiev IF, Shakhidoyatov KhM. 5-Aryl-1,3,4-oxadiazolin-2(3H)-thiones in reactions with of α -haloacetic acid esters. *Chem. Heterocycl. Comp*. 2012, 48:488-491
6. Ismailova DS, Ziyaev AA, Elmuradov BZ, Toshmurodov TT, Bobakulov KhM, Zakirova RP. Targeted synthesis and in vitro bactericidal and fungicidal activities of 2-alkylthio-5-(p-aminophenyl)-1,3,4- oxadiazoles. *Journal of Basic And Applied Research*. 2016, 2:476-479
7. Oxford Diffraction. CrysAlisPro. 2007. Oxford Diffraction Ltd, Abingdon, Oxfordshire, England.
8. Sheldrick GM A short history of SHELX. *Acta Crystallogr. A*. 2008, 64:112-122
9. Tashkhodjaev B, Turgunov KK, Tojiboev AG, Makhmudov US, Ibragimov TF, Saprykina VA, Shakhidoyatov KhM. Crystal solvates and polymorphs of N-(3-methylthio-1,2,4-thiadiazol-5-yl-aminocarbonylmethyl)cytisine. *J. Inclus. Phenom*. 2008, 61:71-76
10. Makhmudov US, Turgunov KK, Tashkhodjaev B, Saprykina VA, Shakhidoyatov KhM. Crystalsolvates of N-(3-ethylthio-1,2,4-thiadiazol-5-yl-aminocarbonylmethyl)cytisine. *Chemistry of Natural Compounds*. 2008, 44:466-471
11. Makhmudov US, Turgunov KK, Tashkhodjaev B, Ibragimov AA, Saprykina VA, Shakhidoyatov KhM. A new polymorphic form and crystal solvates of N-(3-methylthio-1,2,4-thiadiazol-5-yl-aminocarbonylmethyl)cytisine. *J. Inclus. Phenom*. 2010, 66:315-318
12. Makhmudov US, Izotova LYu, Tashkhodjaev B, Ibragimov TF, Shakhidoyatov KhM. 1,2,4-Thiadiazol derivatives of cytosine alkaloid: solvate formation and phase transitions. *J. Inclus. Phenom*. 2012 , 72:375-381
13. Freer AA, Robins DJ, Sheldrake GN. Structures of (-)-cytosine and (-)-N-methylcytosine: tricyclic quinolizidine alkaloids. *Acta Crystallogr. C*. 1987, 43:1119-1122
14. Mascagni P, Christodoulou M, Gibbons WA, Asres K, Phillipson JD, Niccolai N, Mangani S. Solution and crystal structure of cytosine, a quinolizidine alkaloid. *J. Chem. Soc., Perkin Trans. 2*. 1987, 9:1159-1165
15. Barlow RB, Johnson O. Relations between structure and nicotinelike activity: X-ray crystal structure analysis of (-)-cytosine and (-)-lobeline hydrochloride and a comparison with (-)-nicotine-like compounds. *Br. J. Pharmacol*. 1989, 98:799-808
16. Makhmudov US, Rakhimov ShB, Tashkhodzhaev B, Levkovich MG, Vinogradova VI. Molecular structure of N-(*o*- and *p*-hydroxybenzyl)cytisine. *Chemistry of Natural Compounds*. 2008, 44:208-212

17. Abdullaev NP, Makhmudov US, Tashkhodzhaev B, Genzhemuratova G, Levkovich MG, Shakhidoyatov KhM. Structural features of *N*-acylcytisines. *Chemistry of Natural Compounds*. 2009, 45:837-843
18. Estarellas C, Escudero D, Frontera A, Quinonero D, Deya P. *Theor Chem Account*. Theoretical ab initio study of the interplay between hydrogen bonding, cation- π and π - π interactions. 2009. 122: 325-332